

Catalyst Program

Targeting the Relaxin Autocrine Loop in High Grade Serous Ovarian Cancer using RLN2 Neutralizing Monoclonal Antibodies as a Therapeutic Strategy

July 1, 2017 to March 21, 2020

Highlights

- Developing a mAb targeting the Relaxin hormonal pathway as a therapeutic treatment for ovarian cancer
- Establish preclinical support for moving new therapeutic candidates into early-phase clinical testing.

valueted cancer

Ovarian Cancer

The goal of this project is to develop a panel of potent and selective RLN2 neutralizing antibodies that can be tested in high fidelity preclinical models of ovarian cancer.

\$424,500 BioCanRx contribution: \$199,500

RNL2
neutralizing
antibodies

3









About the project

Ovarian cancer has the highest mortality rate of all gynaecological cancers. The standard treatment for patients with ovarian cancer is surgery to reduce tumour burden followed by chemotherapy. Since ovarian cancer is generally not detectable at early stages, most women present with advanced stage disease. Although the majority of women initially respond to chemotherapy, more than 80% of patients will have a recurrence of their disease following treatment, and more than half will die of recurrent disease within 5 years of diagnosis. The identification of specific signaling pathways involved in ovarian tumorigenesis and tumour progression and development of novel treatment strategies to target these pathways is critical. There is an urgent unmet need to identify novel targets that can be drugged therapeutically.

The use of monoclonal antibodies (mAbs) or biologics that block signaling pathways supporting cancer growth has become a major treatment modality for cancer patients in recent years. Dr. Rottapel and his team of researchers propose to develop a mAb targeting the Relaxin hormonal pathway as a therapeutic treatment for ovarian cancer. Relaxin is a peptide hormone expressed in the ovary, which signals through the Relaxin Receptor (RXFP1). They have identified a dependency on Relaxin signaling for the survival and growth of ovarian cancer cells. Blockade of this pathway using a Relaxin targeting mAb or biologic therefore has the potential to antagonize the proliferation of ovarian cancer cells with high specificity.

Using functional genetic shRNA dropout screens the team has uncovered an autocrine loop required for ovarian cancer cell line and xenograft growth involving the mullerian hormone Relaxin and its G-protein coupled receptor RXFP1 (LGR7). They show that recombinant Relaxin stimulates mitogenic and survival signaling pathways and potently induces the expression of VEGF, metalloproteinases (MMPs), and COX2 each known to support epithelial tumourigenicity. Inflammatory cytokines IL-6 and TNFa frequently found in the tumour microenvironment induce Relaxin protein expression in ovarian cancer cell lines further amplifying the Relaxin autocrine loop. Therefore, inhibition of the Relaxin signaling pathway using neutralizing monoclonal antibodies has the potential to block the proliferation and metastatic spread of ovarian cancer cells through multiple mechanisms.



 Test the efficacy of 10 lead mAbs to block proliferation of 4 ovarian cancer cell lines cultured as adherent cells, spheroids and

in xenograft models.

- University Health Network \$128,500

